

氏 名	サトムラ カズヒロ 里村 和浩
所 属	理工学研究科 生命科学専攻
学 位 の 種 類	博士（理学）
学 位 記 番 号	理工博 第 194 号
学位授与の日付	平成 28 年 3 月 25 日
課程・論文の別	学位規則第 4 条第 1 項該当
学位論文題名	減数分裂組換えがアカショウジョウバエの neo-Y 染色体の進化に及ぼす影響（英文）
論文審査委員	主査 教 授 田村 浩一郎 委員 教 授 村上 哲明 委員 准教授 高橋 文

## 【論文の内容の要旨】

本文

The evolution of sex chromosomes is one of crucial issue to understand molecular evolution of eukaryotic species. One hallmark of this process is recombination cessation on Y (W) chromosome evolution. The recombination cessation decreases population size and impairs the efficacy of natural selection of Y chromosome and differentiates sex chromosomes including Y degeneration. Although most of preceding studies showed natural selection under non-recombination environment shaped lack of genetic diversity for Y chromosome, alternative possibility may apply for achiasmatic species, e.g. *Drosophila* species. For example in the case of male achiasmy, Y chromosome has had no chance to introduce genetic diversity through recombination since its origination. Indeed, neo-Y chromosome in *D. miranda* suggested extremely low genetic diversity. Here, in contrast, I focused on a real case of neo-Y chromosome in *D. albomicans* which is in transition from an autosome to a typical Y chromosome but has genetic diversity as rich as gametologous neo-X chromosome.

The neo-X and the neo-Y chromosomes were formed by fusion of sex chromosomes and third chromosomes within 50 million years which is the time *D. albomicans* and its sister species *D. nasuta* diverged. Because most of genes have been maintained intact on the neo-sex chromosomes, I can compare genetic diversity between

gametologous genes. The objective of this study is to clarify the influence of recombination on early stage of Y chromosome evolution in *D. albomicans*.

Using analyses of sequence variations in 53 genes and cross experiments, I elucidated ancestral recombination and current no recombination for the neo-Y chromosome. The time of recombination suppression of the neo-Y chromosome was estimated as a quarter million years ago. My results thus showed the neo-Y chromosome originated from a single chromosomal fusion event, obtained genetic diversity through meiotic recombination during the first few hundred thousand years and diverged from neo-X chromosome after the recombination cessation a quarter million years ago. Consequently, the observed high genetic diversity on the neo-Y chromosome suggested a strong effect of meiotic recombination to introduce genetic variations into the newly arisen sex chromosome.