

Gentle Touch Opens the Gate to the Primary Somatosensory Cortex

2019

Maki Shirato

Contents

1. Abstract.....	4
2. Introduction	6
3. Methods and Materials	12
3.1 Participants.....	12
3.2 Stimuli, trial protocol, and procedure	12
3.3 Analysis of subjective ratings.....	13
3.4 fMRI data analysis.....	13
3.5 Relationships between ROIs and subjective feelings.....	15
3.6 Principal component analysis	15
4. Results	17
4.1 Subjective ratings.....	17
4.2 Differences in brain activity between the C and BF condition	17
4.3 Relationships between ROIs and subjective feelings.....	17
4.4 PCA results	18

4.5 Multiple regression analyses with the right SI, mid-lateral OFC, or ACC as the dependent variable	19
5. Discussion.....	21
6. Table and Figures.....	26
7. References	33
8. Acknowledgments.....	45
9. Research achievements	46

1. Abstract

Objective

Touch is a primary reinforce strongly associated with motivational and affective processes that drive social behavior, and it also plays a critical role in massage therapy. Touch in massage is characterized by gentle touches of the skin involving light pressure effleurage and calm stroking movements intended to increase recipients' pleasure and relaxation. The relationships between basic physical parameters, such as patterns of the hand movements, and their neural bases are important for understanding the effects of gentle touch. However, such studies have not yet been performed. Here, we investigate these relationships and underlying neural mechanisms under two basic movement conditions.

Methods

Using functional magnetic resonance imaging (fMRI), we investigate brain activity induced by circular(C) and back-and-forth (BF) massage of participants' left hands with the experimenter's right hand, ensuring that movements were not unpleasant. We assessed subjective feelings and analyzed fMRI data with principal component analysis (PCA) and correlation analyses to identify associated brain networks.

Results

In C compared with BF, participants felt more positive emotions. There was greater activation of the right primary somatosensory cortex (SI) and left cerebellum (CB), but lower activation of the anterior cingulate cortex (ACC) and periaqueductal gray (PAG) in C compared with BF. There was no significant difference in unpleasant feelings between the conditions. Moreover, co-activation of the left mid-lateral orbitofrontal cortex (OFC), CB, and rostral ventromedial medulla (RVM), and the right SI and posterior insula showed high loadings on Factor 1, which was negatively correlated with unnatural feelings. Meanwhile, co-activation of the ACC and PAG showed high loadings on Factor 2, which was positively correlated with unpleasant feelings.

Conclusion

Our findings suggest somatosensory afferents to the SI are regulated by the descending pain modulatory system under the control of the mid-lateral OFC and ACC, even with mild somatosensory stimulation.

2. Introduction

Touch is a primary reinforce, and one of the foundations of emotion and motivation [1].

Touch associated with positive affect is rewarding, i.e., individuals approach for it, and experience it as pleasant. In contrast, touch associated with negative affect is punishing, i.e., individuals avoid or escape from it, and experience it as unpleasant.

Inter-individual touch is frequently used to communicate positive messages, such as reassurance, comfort, sympathy, and support [2]. Touch from another person can be soothing [3, 4], give rise to pleasurable feelings [5, 6], and potentially suppress pain and negative emotion [7-10]. Skin-to-skin contact between individuals also has a pivotal role in social interactions, sub serving nonverbal communication of intentions and affect.

A romantic caress is a primary expression of affiliative of behavior that reflects the disposition individuals to seek close contact with each other, promoting socio-emotional relationships, pair bonding, and reproduction [11-13]. Therefore, touch is a primary reinforce strongly connected with motivational and affective processes that may drive social behavior [14].

Clinically, massage therapy is often used as an adjunct treatment for various conditions,

including those involving chronic pain and distress [15, 16]. Furthermore, it can enhance growth and cognitive functioning in preterm infants [18-19] and preschoolers [20]. Moreover, massage has beneficial effects in cancer patients by modulating immunity, alleviating depressive symptoms [21], and lessening pain and sleep disturbances [22-24].

Soft touch/stroking, including massage therapy, is characterized by gentle touches of the skin involving light pressure effleurage and calm stroking movements intended to increase the recipient's well-being [25], pleasure [26], and relaxation [27, 28]. Therefore, understanding the influence of physical parameters including the softness, temperature, force, velocity, and movement patterns, is important to clarify their effects and underlying neural mechanisms.

As the human hand is intrinsically soft and warm, softness and temperature may remain within a certain limited range. In addition, force and velocity may also remain within a specific range with the intention to increase the recipient's well-being, pleasure, and relaxation. In contrast, movement patterns (e.g., how to move one's own hand over the back of the recipient's hand) are varied; circular movements may make individuals feel

natural and comfortable, while mechanical movements may make them to feel unnatural and uncomfortable. Accordingly, even a subtle difference in the movement pattern may cause unexpected emotions. Based on these considerations, it is important to clarify the relationships between movement patterns and emotions, and the underlying neural mechanisms. However, such neuroscientific studies have not yet been performed.

The emotional aspects of touch have been explained as positive feelings elicited by activating specific C-tactile (CT) afferents in the skin that signal to areas in the brain involved in positive emotions [29]. Processing of body signals occurs via unmyelinated or thinly myelinated afferents, via interoceptive pathways that signal feelings rather than sensing states, as well as controlling organ functions that do not reach conscious perception. Stroking with a soft brush on hairy skin activates the contralateral posterior insular cortex, as well as the primary somatosensory cortex (SI). The insular cortex is a region of great interest in relation to affective mechanisms, and is considered as a gateway from sensory systems to the emotional systems of the frontal lobe [30, 31].

The medial prefrontal cortex (mPFC)/anterior cingulate cortex (ACC) are also implicated in processing CT-targeted touch [32]. In addition, CTs are tuned to respond to tactile stimuli with the specific thermal characteristics of a gentle caress delivered at

normal skin temperature [33]. Therefore, this reinforces their role in providing a peripheral mechanism to signal pleasant skin-to-skin contact in humans, thereby promoting interpersonal touch and affiliative behavior.

Moreover, an important neural mechanism exists that is known to exert top-down regulation of ascending somatosensory information. Placebo-induced enhancement of pleasant experiences involves upregulation of activity in the posterior insula and SI [34]. In contrast, placebo-induced analgesia involves downregulation of activity in these areas [35-37]. These results indicate that increased early sensory processing of a stimulus of positive valence (e.g., pleasant touch) underpins hyperhedonia, in a similar manner to which reduced processing of an aversive stimulus (e.g., painful touch) underpins analgesia [34]. Placebo-induced improvement of positive and negative hedonic feelings is underpinned by recruitment of common circuitry associated with emotion appraisal. This includes placebo-induced functional coupling between the ventromedial prefrontal cortex (vmPFC) and periaqueductal gray (PAG), which is correlated with increased sensory responses to stroking touch but decreased responses to painful touch [34]. Thus, similar modulatory circuits can either up-or down-regulate early sensory processing depending on whether the expectation is enhancement of

positive or negative hedonic feelings. The PAG–rostral ventromedial medulla (RVM)–spinal cord axis is important for many forms of pro-and anti-nociception in non-human animals, paralleling involvement in human placebo and nocebo effects [38, 39]. This axis is, in turn, governed by the ACC, vmPFC and mid-lateral orbitofrontal cortex (OFC) [36, 40-45], which interact with the PAG–RVM pathway to mediate various pain-modulatory effects as part of the descending pain modulatory system (DPMS) [39].

The above neural mechanism have been investigated by using highly unpleasant tactile stimuli such as those associated with pain, and by using top-down cognitive effects. Thus, whether similar neural mechanisms are activated with mild tactile stimulation such as affiliative and gentle touch remains unconfirmed. A recent fMRI study has shown that there is intrinsic functional connectivity in the DPMS in non-painful situations [46], and another one has shown that self-touch activates the DPMS in a pain-free and stress-free situation [47]. These studies may support the possibility that the DPMS regulates somatosensory afferents based on their emotive valence, even for mild somatosensory stimuli with subtly different movement patterns.

Based on these considerations, we focused in the current study on two basic

movements: circular movement and back-and-forth movement. These movements are fundamental components constituting diverse movement patterns. Circular movement is associated with human and affiliative characteristics that would induce positive feelings, such as gentle, safe, and warm feelings. In contrast, back-and-forth movement has non-human and mechanical characteristics, which would make participants feel unnatural and nervous. We investigated differences in the activity of certain brain regions including the SI, cerebellum (CB), OFC, ACC, insula, and PAG. We conducted analyses to test our hypothesis that the DPMS regulates mild somatosensory afferent signals. We also hypothesized that the OFC activity would be associated with natural and positive emotions induced by circular rubbing, while the ACC activity would be associated with unnatural and nervous feelings induced by back-and-forth rubbing.

3. Methods and Materials

3.1 Participants

We recruited 12 healthy female participants (mean age: 31.5 ± 3.7 years). All participants were right-handed according to the Chapman test (13.3 ± 0.6) and had no history of neurological or psychiatric disorders. Participants provided informed consent to participate in the study. The Research Ethics Committee of Tokyo Metropolitan University approved this study and all methods were performed in accordance with approved guidelines.

3.2 Stimuli, trial protocol, and procedure

Each participant was instructed to relax and close her eyes without thinking about anything specific. We set two experimental conditions, in which one of the authors (TI) who is a professional esthetician, rubbed the back of the participant's hand in a circular motion (C) or rubbed it using a back-and-forth motion (BF). She practiced the rubbing so that she could rub the participant's hand with a maximum force of less than 0.3N, and velocities between 6 cm/sec and 10 cm/sec. Within this range, CT-fibers respond optimally to stimulation [33, 48, 49], and parents naturally stroke their babies [50]. A session consisted of 8 trials (2 conditions \times 4 times), with the trials counterbalanced across participants. A block-design paradigm was applied, with each trial lasting 32 sec

interspersed with rest for 8 sec.

3.3 Analysis of subjective ratings

After the fMRI scans, participants were asked to rate their emotional state while being touched by TI in the same way during the fMRI experiment. We administered nine items to measure subjective feelings: “To what extent did you feel gentle (“gentle”), safe (“safe”), warm (“warm”), comfortable (“comfortable”), preferable (“preferable”), calm (“calm”), unnatural (“unnatural”), nervous (“nervous”), or unpleasant (“unpleasant”)?”. We used visual analog scaling for data collection (0% - 100%). In addition, they were required to rate the perceived intensity on a five-point scale (1 = very weak, 2 = moderately weak, 3 = neutral, 4 = moderately strong, 5 = very strong). Statistical analyses were conducted with SPSS version 21.0 (SPSS, Inc., Chicago, IL).

3.4 fMRI data analysis

Scanning was conducted with a 3.0T MRI system (Achieva Quasar Dual, Philips). BOLD T2*-weighted MR signals were measured with a gradient echo-planar imaging (EPI) sequence (TR = 4,000msec, TE = 35msec, flip angle = 90°, FOV = 23 cm², scan matrix = 128 × 128, total scan time = 324sec, slice thickness = 5mm, 25 slices per volume). Image processing was conducted with Statistical Parametric Mapping 8

(SPM8, Wellcome Department of Imaging Neuroscience, London, UK; <http://www.fil.ion.ucl.ac.uk/spm/software/spm8>). EPI images were realigned and normalized to Montreal Neurological Institute (MNI) stereotactic space. Normalized images were smoothed with an 8 mm full-width half-maximum Gaussian kernel. The data were temporally convolved with the hemodynamic response function (HRF) and high-pass filtered with a cutoff period of 128 sec. Each C and BF condition was modeled with a separate regressor.

We set spherical regions of interest (ROI) in the right SI (40 -31 59) [51], left CB (-36 -61 -29) [52], left mid-lateral OFC (-24 36 -18) [53], right ACC (11 52 15) [54], right posterior insula (42 -31 21) [51], PAG (-6 -33 -18) [55], and RVM (-2 -38 -38) [47].

ROIs with a radius of 5 mm were set for each contrast of C vs. baseline and BF vs. baseline. Subsequently, we tested differences in the eigenvariate values for each ROI between C and BF by using a paired t-test ($P < 0.05$).

Furthermore, we conducted multiple regression analyses with each eigenvariate value for ROIs in the right SI, mid-lateral OFC, and ACC as dependent variable and those of the other ROIs as the independent variables ($P < 0.05$). Multiple regression analyses

were conducted for both C vs. baseline and BF vs. baseline. Moreover, we checked the residuals by performing the Shapiro-Wilks (S-W) test of normality ($P < 0.05$) and calculated the Durbin-Watson (D-W) statistic for the null hypothesis of no autocorrelation.

3.5 Relationships between ROIs and subjective feelings

We conducted multiple regression analyses with each eigenvariate value for the ROIs as the dependent variable and subjective feelings as the independent variables ($P < 0.05$). Moreover, we checked the residuals by performing the S-W test of normality ($P < 0.05$) and calculated the D-W statistic for the null hypothesis of no autocorrelation.

3.6 Principal component analysis

We performed a principal component analysis (PCA) to classify all the ROIs into subgroups by using the Varimax rotation method with Kaiser normalization. In addition, we checked the sampling adequacy by using Bartlett's test of sphericity and the Kaiser-Meyer-Olkin (KMO) test. We also performed multiple regression analyses with scores of each principal component as the dependent variable, and the subjective rating scores as independent variables ($P < 0.05$). We then checked the residuals for all regression analyses by performing the S-W test of normality ($P < 0.05$) and calculated

the D-W statistic for the null hypothesis of no autocorrelation.

Furthermore, to confirm whether simultaneous activation of the RVM and deactivation of the PAG is positively correlated with right SI activity, we performed PCA and calculated the principal component scores for the simultaneous reverse activity and a simple regression analysis between the scores and right SI activity ($P < 0.05$). In addition, we checked the sampling adequacy by using Bartlett's test of sphericity and the Kaiser-Meyer-Olkin test, the residuals for all regression analyses by performing the S-W test of normality ($P < 0.05$), and calculated the D-W statistic for the null hypothesis of no autocorrelation. These statistical analyses were conducted with SPSS version 21.0.

4. Results

4.1 Subjective ratings

Participants showed significantly higher scores for “gentle”, “safe”, “warm”, “comfortable”, “preferable”, and “calm” in C compared with BF. In contrast, they showed significantly higher scores for “unnatural” and “nervous” in BF compared with C. There was no significant difference in “unpleasant” feeling (Figure 1). Moreover, there was also no significant difference in the perceived intensity by Wilcoxon signed-rank test at $P < 0.05$ (C 2.833 ± 0.697 ; BF 2.833 ± 0.333).

4.2 Differences in brain activity between the C and BF condition

The right SI ($t = 3.077$, $P = 0.011$) and left CB ($t = 3.587$, $P = 0.004$) showed significantly higher activity in C compared with BF. In contrast, the ACC ($t = -2.248$, $P = 0.0461$) and PAG ($t = -2.492$, $P = 0.029$) showed significantly higher activity in BF compared with C (Figure 2).

4.3 Relationships between ROIs and subjective feelings

Activity in the left CB, right posterior insula, and PAG were positively correlated with feeling “safe”(CB: $t = 2.380$, $P = 0.026$, adjusted $R^2 = 0.169$; S-W statistic = 0.949, $P = 0.255$; D-W statistic = 2.578; posterior insula: $t = 3.233$, $P = 0.004$, adjusted $R^2 = 0.291$; S-W statistic = 0.919, $P = 0.054$; D-W statistic = 1.783), and “unpleasant”(PAG:

$t=2.825$, $P=0.010$, adjusted $R^2=0.233$; S-W statistic = 0.972, $P=0.720$; D-W statistic = 1.829) (Figure 3).

In contrast, activity in the right SI, RVM, mid-lateral OFC, and ACC was negatively correlated with feeling “unnatural”(SI: $t=-3.726$, $P=0.001$, adjusted $R^2=0.359$; S-W statistic = 0.935, $P=0.126$; D-W statistic = 2.012; RVM: $t=-2.214$, $P=0.037$, adjusted $R^2=0.145$; S-W statistic = 0.950, $P=0.272$; D-W statistic = 1.797), “unpleasant”(OFC: $t=-2.835$, $P=0.010$, adjusted $R^2=0.234$; S-W statistic = 0.924, $P=0.070$; D-W statistic = 1.924), and “comfortable”(ACC: $t=-2.395$, $P=0.026$, adjusted $R^2=0.171$; S-W statistic = 0.954, $P=0.329$; D-W statistic = 1.633) (Figure 3).

4.4 PCA results

We extracted two principal components: 1) brain activity associated with positive emotion (48.544% of the total variance); and 2) brain activity associated with negative emotion (21.902% of the total variance). A subsequent exploratory factor analysis including seven predictors for the ROI activity scores revealed that these predictors converged on two factors (Figure 4). The first (Factor 1) included activity in the SI, CB, posterior insula, mid-lateral OFC, and RVM. The second (Factor 2) included activity in the ACC and PAG (Table, Figure 4).

We confirmed the sampling adequacy with Bartlett's test of sphericity ($\chi^2=89.556$, $df=21$, $P<0.0001$) and KMO's test (KMO measure=0.627). Moreover, the score for Factor 1 was negatively correlated with “unnatural” feelings ($t=-2.870$, $P=0.009$; adjusted $R^2=0.239$; S-W statistic = 0.982, $P=0.936$; D-W statistic = 2.132). In addition, the score for Factor 2 showed was significantly positively correlated with “unpleasant” feelings ($t=2.478$, $P=0.021$; adjusted $R^2=0.183$; S-W statistic = 0.980, $P=0.900$; D-W statistic = 1.366). Furthermore, there was a significant positive correlation between the score for simultaneous activation of the RVM and deactivation of the PAG (55.954% of the total variance) and right SI activity ($t=4.361$, $P<0.0001$, adjusted $R^2=0.439$; S-W statistic = 0.940, $P=0.165$; D-W statistic = 2.705) (Figure 5).

4.5 Multiple regression analyses with the right SI, mid-lateral OFC, or ACC as the dependent variable

Activity in the right SI was positively correlated with that in the left CB ($t=2.199$, $P=0.040$), RVM ($t=2.783$, $P=0.012$), and right posterior insula ($t=2.561$, $P=0.019$). In contrast, activity in the right SI was negatively correlated with that in the PAG ($t=-2.709$, $P=0.014$; adjusted $R^2=0.761$; S-W statistic = 0.393, $P=0.393$; D-W statistic = 1.707) (Figure 6). Moreover, activity in the mid-lateral OFC was positively correlated

with that in the RVM ($t= 5.327$, $P<0.0001$), and negatively correlated with that in the PAG ($t= -2.375$, $P= 0.027$; adjusted $R^2= 0.607$; S-W statistic $=0.742$, $P= 0.742$; D-W statistic $=1.540$)(Figure 6). Activity in the ACC was positively correlated with that in the PAG ($t= 2.581$, $P= 0.017$; adjusted $R^2= 0.197$; S-W statistic $= 0.928$, $P= 0.088$; D-W statistic $=2.873$)(Figure 6).

5. Discussion

In the present study, participants reported that they felt more positive feelings when circularly rubbed on the back of the left hand compared with being rubbed back and forth on the same hand. In contrast, they reported that they felt more unnatural and nervous when rubbed back and forth compared with being circularly rubbed. Moreover, there was no significant difference in feelings of unpleasantness. As back-and-forth linear movements have non-human and mechanical characteristics, they might make participants feel unnatural and nervous. In contrast, circular movements have human and affiliative characteristics, which might make participants experience positive feelings.

Furthermore, in BF compared with C, there was significantly more activation in the ACC and PAG, but significantly less in the right SI and left CB. Furthermore, ACC activity was negatively correlated with “comfortable” feelings and PAG activity was positively correlated with “unpleasant” feelings. Meanwhile, right SI activity was negatively correlated with “unnatural” feelings and left CB activity was positively correlated with “safe” feelings. These results suggest that ACC and PAG activity is related to negative emotions associated with being rubbed, while right SI and left CB

activity are related to the positive emotions evoked by rubbing.

The ACC receives somatosensory information via the insula and OFC, which are both well connected with the ACC [56]. ACC activation was observed to increase following a suggested increase in subjective unpleasantness, but to decrease following a suggested decrease in subjective unpleasantness [57]. This indicates that the ACC is involved in representing affective qualities of a stimulus rather than its physical intensity. In fact, there was no significant difference in the perceived intensity between C and BF.

Activity in the ACC is also positively correlated with unpleasantness of pain in healthy individuals [58], indicating that this region is particularly involved in affective responses to pain. Moreover, the ACC controls the brainstem, including the PAG and RVM [59, 60], during both opioid and placebo analgesia [39, 61]. The ACC has one of the highest levels of opioid receptor binding in the cortex [62], and positron emission tomography (PET) studies indicate that the binding potential is specifically highest in the ACC [63, 64].

The PAG plays a pivotal role in the DPMS [65]. It is functionally connected with the ACC [46] and participates in the descending modulation of pain. Indeed, our results that

ACC activity is positively correlated with PAG activity, which in turn was negatively correlated with right SI activity, suggests involvement of the DPMS. Moreover, the DPMS can function even in the absence of painful/stressful stimulation, because intrinsic functional connectivity among DPMS regions (including the ACC and RVM) has been demonstrated in such situations [46]. Moreover, the DPMS is activated during self-touching behaviors in pain-free and stress-free situations [47].

A positive correlation has been observed between behavioral opioid analgesia and opioid-induced suppression of neuronal responses to noxious stimuli in the RVM [66]. The RVM plays a critical role in both inhibition and facilitation of pain through interactions with the spinal cord. Both off-and on-cells in the RVM project to the spinal dorsal horn, indicating that they exert modulatory influences on nociceptive/non-nociceptive inputs [67]. RVM on-cells are directly inhibited by opioids, and it is suggested that these cells express mu-opioid receptors [68]. Moreover, increased RVM neuronal responses to noxious stimuli observed in human imaging studies indicate on-cell activity [69, 70]. In fact, we observed a positive correlation between RVM activity and right SI activity in the present study. Furthermore, simultaneous activation of the RVM and deactivation of the PAG was positively

correlated with right SI activity. This suggests that PAG activity inhibits on-cell activity in the RVM, which suppresses right SI activity. In fact, PAG neurons modulate nociception by directly inhibiting on-cells in the RVM via GABAergic neurons [71]. Conversely, somatosensory afferent signals were enhanced (facilitated) by circular rubbing compared with back-and-forth rubbing. We suggest that the mid-lateral OFC downregulates PAG activity, which inhibits RVM activity and/or directly upregulates RVM activity, based on reward/saliency evaluation processes in the mid-lateral OFC. This suggestion is supported by results indicating that mid-lateral OFC activity is negatively correlated with PAG activity, and positively correlated with RVM activity. Moreover, the mid-lateral OFC showed a negative correlation with “unpleasant” feelings. Furthermore, co-activation of the left mid-lateral OFC, CB, and RVM, and of the right SI and posterior insula showed high factor loading on Factor1, which was negatively correlated with “unnatural” feelings. However, these co-activations showed low factor loading on Factor 2, which was positively correlated with “unpleasant” feelings. The posterior and mid-insula receive somatosensory information through projections from the SI [72, 73] and directly from the thalamus [74]. The OFC receives somatosensory information from the granular insula and directly from the SI [75-77]. As such, the OFC can mediate integration of such information and evaluate its

rewarding (salient) characteristics. Indeed, a meta-analysis has suggested that sensory pleasure is most faithfully represented within a mid-lateral OFC site [78].

Neuroimaging studies of placebo effects on physical pain suggest that activation of the ACC, OFC, and vmPFC influences a in by activating the descending pain regulatory pathways in the brainstem, especially opioidergic mechanisms in the PAG [39, 45, 79-83]. The DPMS is important for many forms of pro-and anti-nociception in non-human animals, paralleling involvement in human placebo and nocebo effects. The present findings show that somatosensory afferents to the SI are regulated by the DPMS under the control of the mid-lateral OFC and ACC, even under mild somatosensory stimulation situations.

6. Table and Figures

Figure 1. Differences in subjective feelings between C and BF.




Figure 2. Relationships between ROIs and subjective feelings.




Figure 3. Correlations between brain activity and subjective feelings.




Table 1. Exploratory factor structure of brain activity

Factor analysis	Factor 1		Factor 2
Brain regions	Factor loading	Factor loading	
RVM	0.926	0.157	
R SI	0.906	-0.204	
L CB	0.801	0.023	
Mid-lateral OFC	0.787	-0.245	
R posterior insula	0.546	0.056	
ACC	0.109	0.734	
PAG	-0.190	0.924	
Regression analysis	feeling	unnatural	unpleasant
	correlation	negative (-)	positive (+)

Figure 4.2D scatter plot of principal components (PCs) of brain activity. The x-axis and the y-axis represent PC1 and PC2, respectively.




Figure 5. Significant positive correlation between the score for simultaneous activation of the RVM and deactivation of the PAG and right SI activity.





Figure 6. Multi-regression analyses with the right SI, mid-lateral OFC, or ACC as the dependent variable. Each arrow represents the correlation, and the tail and the head represent an independent variable and the dependent variable, respectively.



References

1. Rolls ET. *Emotion Explained*. Oxford University Press, Oxford (2005).
2. Hertenstein MJ, Verkamp JM, Kerestes AM, et al. The communicative functions of touch in humans, nonhuman primates, and rats: a review and synthesis of the empirical research. *Genet. Soc. Gen. Psychol. Monogr* 132 (1), 5-94 (2006).
3. Feldman R, Singer M, Zagoory O. Touch attenuates infants' physiological reactivity to stress. *Dev. Sci* 13 (1), 271–278 (2010).
4. Fairhurst MT, Löken L, Grossmann T. Physiological and behavioral responses reveal 9-month-old infants' sensitivity to pleasant touch. *Psychol. Sci* 25(5), 1124-1131 (2014).
5. Löken LS, Wessberg J, Morrison I, et al. Coding of pleasant touch by unmyelinated afferents in humans. *Nat. Neurosci* 12, 547–548 (2009).
6. Morrison I, Löken LS, Olausson H. The skin as a social organ. *Exp Brain Res* 204(3): 305-314 (2010).
7. Coan JA, Schaefer HS, Davidson RJ. Lending a hand: social regulation of the neural response to threat. *Psychol.Sci* 17(1), 1032–1039 (2006).
8. Liljencrantz J, Ellingsen DM, Leknes S, et al. "C-tactile afferent stimulation modulate pain perception," in Society for Neuroscience, New Orleans, LA. (2012).

9. Mancini F, Nash T, Iannetti GD, et al. Pain relief by touch: a quantitative approach. *Pain* 155(1), 635–642 (2014).
10. Mancini F, Beaumont AL, Hu L, et al. Touch inhibits subcortical and cortical nociceptive responses. *Pain* 156, 1936–1944 (2015).
11. Dunbar RI. The social role of touch in humans and primates: behavioural function and neurobiological mechanisms. *Neurosci. Biobehav. Rev* 34(1), 260–268 (2010).
12. Gallace A, Spence C. The science of interpersonal touch: an overview. *Neurosci. Biobehav. Rev* 34(1), 246–259 (2010).
13. Morrison I, Löken LS, Olausson H. The skin as a social organ. *Exp. Brain. Res* 204(1), 305–314 (2010).
14. Rolls ET. The affective and cognitive processing of touch, oral texture, and temperature in the brain. *Neurosci. Biobehav. Rev* 34(1), 237–245 (2010).
15. Sherman KJ, Cherkin DC, Hawkes RJ, et al. Randomized trial of therapeutic massage for chronic neck pain. *The Clinical Journal of Pain*, 25(3)233–238 (2009).
16. Field T. Massage therapy. *Medical Clin. Am* 86(1), 163–171 (2002).
17. Field T, Diego M, Hernandez-Reif M, et al. Insulin and insulin-like growth factor I (IGF-1) increase in preterm infants following massage therapy. *J. Dev. Behav. Pediatr* 29(6), 463–466 (2008).

18. Field T, Diego M, Hernandez-Reif M. Preterm infant massage therapy research: a review. *Infant Behavi. Dev* 33(2), 115–124 (2010).
19. Procianoy RS, Mendes EW, Silveira RC. Massage therapy improves neurodevelopment outcome at two years corrected age for very low birth weight infants. *Early. Human. Dev* 86(1), 7–11 (2010).
20. Hart S, Field T, Hernandez-Reif M, et al. Preschoolers' cognitive performance improves following massage. *Early. Child. Dev. Care* 143(1), 59–64 (1998).
21. Hou WH, Chiang PT, Hsu TY, et al. Treatment effects of massage therapy in depressed people: a meta-analysis. *J. Clin. Psychiatry* 71(7), 894–901 (2010).
22. Russell NC, Sumler SS, Beinhorn CM, et al. Role of massage therapy in cancer care. *J. Altern. Complement. Med* 14(2), 209–214 (2008).
23. Hughes D, Ladas E, Rooney D, et al. Massage therapy as a supportive care intervention for children with cancer. *Oncol. Nurs. Forum* 35(3), 431–442 (2008).
24. Calenda E. Massage therapy for cancer pain. *Curr. Pain. Headache. Rep* 10(4), 270–274 (2006).
25. Cronfalk BS, Strang P, Ternestedt BM. Inner power, physical strength and existential well-being in daily life: relatives' experiences of receiving soft tissue massage in palliative home care. *J. Clin. Nurs* 18(1), 2225-2233 (2009).

26. Bergsten U, Petersson IF, Arvidsson B. Perception of tactile massage as a complement to other forms of pain relief in rheumatic disease. *Musculoskeletal Care* 3(1), 157-167 (2005).
27. Billhult A, Maatta S. Light pressure massage for patients with severe anxiety. *Complement. Ther. Clin. Pract* 15, 96-101 (2009).
28. Goodfellow LM. The effects of therapeutic back massage on psychophysiologic variables and immune function in spouses of patients with cancer. *Nurs. Res* 52, 318-328 (2003).
29. Olausson H, Cole J, Rylander K, et al. Functional role of unmyelinated tactile afferents in human hairy skin: sympathetic response and perceptual localization. *Exp. Brain Res* 184, 135–140 (2008).
30. Augustine JR. Circuitry and functional aspects of the insular lobe in primates including humans. *Brain Res. Rev* 22(3), 229-244 (1996).
31. Craig AD. Interoception and emotion. In *Handbook of Emotions*, Third Edition, Lewis M, Haviland-Jones JM, Barrett LF eds. 272–288, Guilford Publications, NY (2008).
32. Gordon I, Voos AC, Bennett RH, et al. Brain mechanisms for processing affective touch. *Hum. Brain. Mapp* 34, 914–922 (2013).

33. Ackerley R, Backlund Wasling H, Liljencrantz J, et al. Human C-tactile afferents are tuned to the temperature of a skin-stroking caress. *J. Neurosci* 34, 2879–2883 (2014).
34. Ellingsen D-M, Wessberg J, Eikemo M, et al. Placebo improves pleasure and pain through opposite modulation of sensory processing. *Proceedings of the National Academy of Sciences of the United States of America* 110(44), 17993-17998 (2013).
35. Amanzio M, Benedetti F, Porro CA, et al. Activation likelihood estimation meta-analysis of brain correlates of placebo analgesia in human experimental pain. *Hum. Brain. Mapp* 34(3), 738–752 (2013).
36. Eippert F, Bingel U, Schoell ED, et al. Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron* 63(4), 533–543 (2009).
37. Lu HC, Hsieh JC, Lu CL et al. Neuronal correlates in the modulation of placebo analgesia in experimentally-induced esophageal pain: A 3T-fMRI study. *Pain* 148(1), 75–83 (2010).
38. Watkins LR, Mayer DJ. Organization of endogenous opiate and non-opiate pain control systems. *Science* 216, 1185–1192 (1982).
39. Wager TD, Atlas LY. The neuroscience of placebo effects: connecting context,

- learning and health. *Nat. Rev. Neurosci* 16, 403–418 (2015).
40. Atlas LY, Whittington RA, Lindquist MA, et al. Dissociable influences of opiates and expectations on pain. *J. Neurosci* 32, 8053–8064 (2012).
 41. Wiech, K. Lin CS, Brodersen KH et al. Anterior insula integrates information about salience into perceptual decisions about pain. *J. Neurosci.* 30, 16324–16331 (2010).
 42. Lee HF, Hsieh JC, Lu CL, et al. Enhanced affect/cognition-related brain responses during visceral placebo analgesia in irritable bowel syndrome patients. *Pain* 153, 1301–1310 (2012).
 43. Atlas LY, Bolger N, Lindquist MA, et al. Brain mediators of predictive cue effects on perceived pain. *J. Neurosci* 30, 12964–12977 (2010).
 44. Kong J, Jensen K, Loiotile R, et al. Functional connectivity of the frontoparietal network predicts cognitive modulation of pain. *Pain* 154, 459–467 (2013).
 45. Petrovic P, Kalso E, Petersson KM, et al. Placebo and opioid analgesia — imaging a shared neuronal network. *Science* 295, 1737–1740 (2002).
 46. Kong J, Tu PC, Zyloney C, et al. Intrinsicfunctional connectivity of the periaqueductal gray, a resting fMRI study. *Behav. Brain. Res* 211(2), 215–219 (2010).
 47. Kikuchi Y, Shirato M, Machida A, et al. The neural basis of self-touch in a pain-free

- situation. *Neuropsychiatry (London)* 8(1), 186–196 (2018).
48. Olausson H, Wessberg J, Morrison I, et al. The neurophysiology of unmyelinated tactile afferents. *Neurosci. Biobehav. Rev* 34 (2), 185–191 (2010).
 49. Löken LS, Wessberg J, Morrison I, et al. Coding of pleasant touch by unmyelinated afferents in humans, *Nat. Neurosci* 12 (5), 547–548 (2009).
 50. Croy A, Luong C, Tricoli E, et al. Interpersonal stroking touch is targeted to C tactile afferent activation, *Behav. Brain. Res* 297, 37-40 (2016).
 51. Ackerley R, Hassan E, Curran A, et al. An fMRI study on cortical responses during active self-touch and passive touch from others. *Front. Behav. Neurosci* 6:51 (2012).
 52. Gordon I, Voos AC, Bennett RH, et al. Brain mechanisms for processing affective touch. *Hum. Brain. Mapp* 34, 914-922 (2013).
 53. Hardin MG, Pine DS, Ernst M. The influence of context valence in the neural coding of monetary outcomes. *Neuroimage* 48, 249-257 (2009).
 54. Zeidan F, Grant JA, Brown BA, et al. Mindfulness meditation-related pain relief: evidence for unique brain mechanisms in the regulation of pain. *Neurosci. Letters* 520, 165-173 (2012).
 55. Wiech K, Edwards R, Moseley GL, et al. Dissociable neural mechanisms underlying the modulation of pain and anxiety? an fMRI pilot study. *PLOS. ONE* 9(12) (2014).

56. Van Hoesen GW, Morecraft RJ, Vogt BA. Connections of the monkey cingulate cortex. In: The neurobiology of the cingulate cortex and limbic thalamus: a comprehensive handbook, Vogt BA, Gabriel Meds, 249–284 Birkhäuser, Boston, MA (1993).
57. Rolls ET, O'Doherty J, Kringelbach ML, et al. Representations of pleasant and painful touch in the human orbitofrontal and cingulate cortices. *Cereb. Cortex* 13(3), 308-317 (2003).
58. Brown CA, Jones AK. Meditation experience predicts less negative appraisal of pain: electrophysiological evidence for the involvement of anticipatory neural response. *Pain* 150, 428-438 (2010).
59. Fields H. State-dependent opioid control of pain. *Nat. Rev. Neurosci* 5(7), 565-575 (2004).
60. Petrovic P. Opioid and placebo analgesia share the same network. *Semin. Pain. Med* 3(1), 31-36 (2005).
61. Zubieta JK, Ketter TA, Bueller JA, et al. Regulation of human affective responses by anterior cingulate and limbic mu-opioid neurotransmission. *Arch. Gen. Psychiatry* 60(11), 1145-1153 (2003).
62. Vogt BA, Wiley RG, Jensen EL. Localization of Mu and delta opioid receptors to

- anterior cingulate afferents and projection neurons and input/output model of Mu regulation. *Exp. Neurol* 135(2), 83-92 (1995).
63. Jones AK, Qi LY, Fujirawa T, et al. In vivo distribution of opioid receptors in man in relation to the cortical projections of the medial and lateral pain systems measured with positron emission tomography. *Neurosci. Lett* 126(1), 25-28 (1991).
64. Willoch F, Schindler F, Wester HJ, et al. Central poststroke pain and reduced opioid receptor binding within pain processing circuitries: a [¹¹C] diprenorphine PET study. *Pain* 108(3), 213-220 (2004).
65. Basbaum AI, Fields HL. Endogenous pain control systems: brainstem spinal pathways, and endorphin circuitry. *Annu. Rev. Neurosci* 7, 309-338(1984).
66. Hopkins E, Spinella M, Pavlovic ZW, et al. Alterations in swim stress-induced analgesia and hypothermia following serotonergic or NMDA antagonists in the rostral ventromedial medulla of rats. *Physiol. Behav* 64(3), 219-225(1998).
67. Fields HL, Malick A, Burstein R. Dorsal horn projection targets of ON and OFF cells in the rostral ventromedial medulla. *J. Neurophysiol* 74(4), 1742-1759 (1995).
68. Heinricher MM, Morgan MM, Fields HL. Direct and indirect actions of morphine on medullary neurons that modulate nociception. *Neuroscience* 48(3), 533-543 (1992).

69. Dunckley P, Wise RG, Fairhurst M, et al. A comparison of visceral and somatic pain processing in the human brainstem using functional magnetic resonance imaging. *J. Neurosci* 25(32), 7333-7341 (2005).
70. Gwilym SE, Keltner JR, Warnaby CE, et al. Psychophysical and functional imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis. *Arthritis. Rheum* 61(9), 1226-1234 (2009).
71. Morgan MM, Whittier KL, Hegarty DM, et al. Periaqueductal Gray neurons project to spinally projecting GABAergic neurons in the rostral ventromedial medulla. *Pain* 140(2), (2008).
72. Mufson EJ, Mesulam MM. Insula of the old world monkey: II: afferent cortical input and comments on the claustrum. *J. Comp. Neurol* 212, 23–37 (1982).
73. Friedman DP, Murray EA, O'Neill JB, et al. Cortical connections of the somatosensory fields of the lateral sulcus of macaques: evidence for a corticolimbic pathway for touch. *J. Comp. Neurol* 252, 23–347 (1986).
74. Craig AD, Bushnell MC, Zhang ET, et al. A thalamic nucleus specific for pain and temperature sensation. *Nature* 372, 770–773 (1994).
75. Barbas H. Anatomic organization of basoventral and mediodorsal visual recipient prefrontal regions in the rhesus monkey. *J. Comp. Neurol* 276, 313–342 (1988).

76. Morecraft RJ, Geula C, Mesulam MM. Cytoarchitecture and neural afferents of orbitofrontal cortex in the brain of the monkey. *J. Comp.Neurol* 232, 341–358 (1992).
77. Carmichael ST, Price JL. Sensory and premotor connections of the orbital and medial prefrontal cortex of macaque monkeys. *J. Comp.Neurol* 363, 642–664 (1995).
78. Kringelbach ML, Rolls ET. The functional neuroanatomy of the human orbitofrontal cortex: evidence from neuroimaging and neuropsychology. *Prog.Neurobiol* 72, 341-372 (2004).
79. Vogt BA, Sikes RW, Vogt LJ. Anterior cingulate cortex and the medial pain system. In: *Neurobiology of cingulate cortex and limbic thalamus*, 313–344, Springer, NY (1993).
80. Benedetti F, Arduino C, Amanzio M. Somatotopic activation of opioid systems by target-directed expectations of analgesia. *J. Neurosci* 19, 3639–3648 (1999).
81. Bingel U, Schoell E, Büchel C. Imaging pain modulation in health and disease. *Curr. Opin. Neurol* 20, 424–431 (2007).
82. Scott DJ, Stohler CS, Egnatuk CM, et al. Placebo and nocebo effects are defined by opposite opioid and dopaminergic responses. *Arch. Gen. Psychiatry* 65, 220 –231

(2008).

83. Eippert F, Bingel U, Schoell ED, et al. Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron* 63, 533–543 (2009).

7. Acknowledgments

I would like to express my sincerest gratitude to my supervisor, Professor Yoshiaki Kikuchi, for providing me with this amazing opportunity as a Ph. D student in his laboratory.

I would also like to express my deepest appreciation to Dr. Madoka Noriuchi for her close support and guidance, her encouragement and all the invaluable discussions that both made my research a success and my study life unforgettable.

I am very grateful to Dr. Akiko Machida and Ms. Toshie Inoue for their valuable cooperation in my experiments, and to Professor Sadaki Takata for supporting me.

Finally, I would like to extend my love and thanks to my husband Satoshi and my children Leo and Mao for their endless love, understanding, support, encouragement and sacrifices throughout my study.

December 20, 2019

Maki Shirato

8. Research achievements

Publications

Academic journal published papers

1. Gentle Touch Opens the Gate to the Primary Somatosensory Cortex.

Shirato M., Kikuchi Y., Machida A., Inoue T., Noriuchi M. Neuropsychiatry

(London) (2018) 8(5), 1696-1707, 2018

2. The Neural Basis of Self-Touch in a Pain-Free Situation.

Kikuchi Y., Shirato M., Machida A., Inoue T., Noriuchi M. Neuropsychiatry

(London) (2018) 8(1), 186-196, 2018

3. Effects of Cosmetic Therapy on Cognitive Function in Elderly Women Evaluated by Time-Resolved Spectroscopy Study.

Machida A., Shirato M., Tanida M., Kanemaru C., Nagai S., Sakatani K. Adv Exp

Med Biol. 876, 289-295, 2016

4. 化粧療法が高齢者の脳波にもたらす変化.

町田明子, 上田泰士, 渥美圭大, 白土真紀, 高田定樹. 老年精神医学雑誌, 24,

9, 915-927, 2013

5. 脳波を用いた高齢者に対する化粧療法の効果検証.

町田明子, 白土真紀, 高田定樹, 八木透. 老年精神医学雑誌, 23, 8,

978-987, 2012

6. 軽度認知症高齢者に対するハンドマッサージの効果検証.

白土真紀, 井上俊枝, 石川好江, 町田明子, 高田定樹, 安藤弥生, 工藤千秋.

日本早期認知症学会誌, 5, 1, 22-27, 2012

Other Publications

1. 『エビデンスに基づく認知症補完療法へのアプローチ』.

長谷川和夫・武者利光監, 工藤千秋 編/白土真紀, 高田定樹, 井上俊枝 著.

株式会社ぱーそん書房, p58-62, 2012

2. 50代からさらに美しく エイジング美容.

青木昭子, 井上俊枝, 上田美江子, 白土真紀 (資生堂美の匠チーム). 株式会

社ポプラ社, 2011

International conferences

1. Effects of Cosmetic Therapy on Cognitive Function in Elderly Women Evaluated by

Time-Resolved Spectroscopy Study.

Sakatani K., Machida A., Shirato M., Kanemaru C., Nagai S. ISOTT2014, UCL,

London, UK, Abstract, p.100, 2014

2. Physical and mental effects of cosmetic therapy for the elderly.

Shirato M., Machida A., Kanemaru C., Nagai S., Takata S. The International Federation of Societies of Cosmetic Chemists (IFSCC2012) Abstract, 43-44, Proceeding No. 036, 2012

3. Evaluation of the effect of cosmetic therapy by EEG.

Machida A., Shirato M., Takata S, Atsumi Y., and Yagi T. Proceedings of the 2011 Biomedical Engineering International Conference (BNEiCON-2011),48-53, 2011

4. Effects of cosmetic therapy on prefrontal cortex activity and salivary cortisol level.

Machida A., Shirato M., Kanemaru C., Kio K., Nagai S., Takata S., Ikeda M. and Sakatani K. 29th International Congress of Clinical Neurophysiology (ICCN) 2010, Clinical Neurophysiology Vol. 121 Suppl.1 OCTOBER 2010, S206, 2010

Domestic conferences

1. 歩行の審美性に関する運動学的考察.

水村（久埜）真由美, 河野由, 白土真紀, 平尾直靖, 根岸茜子. 第39回バイオメカニズム学術講演会, 筑波, 講演予稿集, p2-3, 2018

2. スキンケア化粧品の愛用意識に関するfMRIを用いた認知神経科学的研究.

磯部寛子, 則内まどか, 白土真紀, 瀬戸山三佳, 八巻智史, 平尾直靖, 菊池吉晃. 第43回日本化粧品学会, 東京, 講演要旨, p27, 2018

3. アイメイクによる黒目の過大錯視.

松下戦具, 今藤咲智, 森川和則, 白土真紀, 山南春奈. 日本顔学会学第22回大会, 大阪, 日本顔学会誌, No1, 82, 2017

4. 顔の要素間の大きさ同化錯視 ―顔全体処理と錯視の関連―.

孫卉雯, 森川和則, 松下戦具, 白土真紀, 山南春奈. 日本心理学会第81回大会, 福岡, 2017

5. 眉と鼻筋の化粧による小顔錯視効果の測定

松下戦具, 森川和則, 管万純, 長島愛, 白土真紀. 日本顔学会学第21回大会, 東京, 日本顔学会誌, No1, 46, 2016

6. 二重まぶたで目が大きく見える錯視効果

松下戦具, 杉澤みなみ, 森川和則, 白土真紀. 日本認知心理学会学第13回大会, 東京, 2015

7. メーキャップによる自己顔印象変化の脳イメージング研究.

白土真紀, 町田明子, 礪部寛子, 則内まどか, 菊池吉晃. 日本顔学会学第19回大会, 東京, 日本顔学会誌, No1, 46, 2014

8. 香りによる懐かしさの想起 ―黒質/海馬の機能的結合―.

白土真紀, 大場健太郎, 森下薫, 則内まどか, 小早川達, 菊池吉晃. 日本生理人類学会第70回大会, 九州, 2014

9. メーキャップによる自己顔変化についての認知神経科学的研究.

白土真紀, 町田明子, 磯部寛子, 則内まどか, 菊池吉晃. 第 75 回 SCCJ 研究討論会, 東京, 2014

10. 触れる行為の認知神経科学的検討.

白土真紀, 町田明子, 井上俊枝, 高田定樹, 則内まどか, 菊池吉晃. 日本生理人類学会第 67 回大会, 東京, 日本生理人類学会誌, Vol.12, 特別号(2), 第 67 回大会要旨集, 86-87, 2012

11. 脳科学的手法を用いた製品や店頭サービスの開発.

白土真紀, 高田定樹. 日本生理人類学会, 第 67 回大会, 東京, 日本生理人類学会誌, Vol.12, 特別号(2), 第 67 回大会要旨集, 26-27, 2012

12. 化粧療法による高齢者への心身効果の検証.

白土真紀, 町田明子, 金丸千枝美, 永井幸恵, 高田 定樹. IFSCC 2012 ヨハネスブルグ大会・国内報告会, 東京, 講演要旨集, 19, 42, 2012

13. ハンドマッサージの影響についての生理心理学的検討ー触れる行為の認知神経科学的検討とあわせてー.

白土真紀, 町田明子, 井上俊枝, 高田 定樹, 則内まどか, 菊池吉晃. 第 75 回 SCCJ 研究討論会, 東京, 2012, 講演要旨集, 20-21, 52, 2012

14. 認知症の改善を目的とした高齢者への化粧セラピーの定量評価に関する研究.

渥美圭大, 町田明子, 白土真紀, 高田定樹, 八木透. 電気学会医用・生体工学研究会, 東京, 2011

15. 認知症の改善を目的とした高齢者への化粧セラピーの定量評価に関する研究.

渥美圭大, 町田明子, 白土真紀, 高田定樹, 八木透. 第 26 回生体・生理工学シンポジウム, 草津, 2011

16. 脳波を用いた化粧シュミレーターの効果の検証に関する研究.

山本晃嗣, 町田明子, 白土真紀, 高田定樹, 八木透. 電気学会医用・生体工学研究会, 東京, 2011

17. 化粧の視覚的效果が脳波に及ぼす研究.

山本晃嗣, 町田明子, 白土真紀, 高田定樹, 八木透. 電気学会電子・情報・システム部門大会講演論文集, 東京, 2011

18. DIMENSION・NAT を用いた高齢者化粧療法の効果検証.

町田明子, 白土真紀, 高田定樹, 八木透. 第 3 回臨床脳電位研究会学術大会, 東京, 2011